
Management of open apex in maxillary incisors with mta - a retrospective study

PRIADARSINI T¹, SOWMYA K^{2*}, DHANRAJ GANAPATHY³

¹Saveetha dental college and hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University Chennai - 77

²Senior lecturer, Department of Conservative Dentistry and Endodontics, Saveetha Dental College and Hospitals Saveetha Institute of Medical and Technical Sciences Saveetha University, Chennai-77

³Professor, Department of Prosthodontics, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences Saveetha University, Chennai-77

*Corresponding Author

Email ID: 151501017.sdc@saveetha.com¹, sowmyak.sdc@saveetha.com, dhanraj.sdc@saveetha.com

Abstract: Open apices commonly arise secondary to pulpal necrosis as a result of caries or trauma in an immature tooth with incomplete root formation. The conventional method of apexification with calcium hydroxide has certain disadvantages such as a very long period of treatment, tooth fracture and an incomplete calcification of the bridge. MTA has gained importance as an alternative treatment for management of open apices as it overcomes these disadvantages. The aim of this study was to determine the number, age and gender distribution of MTA apexification done in maxillary incisors. This study included data of 41 patients who had undergone apexification using MTA in maxillary incisors. The data was collected from the case records of patients visiting Saveetha dental College for treatment. A total of 47 teeth that underwent MTA apexification were identified. Of the total subjects, 83% were males and 17% were females. 51.06% of the teeth that underwent MTA apexification belonged to the age group <15 years and 48.04% to the group >15 years. Statistical analysis was done using IBM SPSS software version 20.0. No significant association was found between gender and different age groups in patients that underwent MTA apexification (P value= 0.701 >0.05; Fisher's exact test). MTA apexification was done predominantly in males in both age groups.

Keywords: Apexification; Incisors; MTA; Open apex; Trauma innovative technique

INTRODUCTION

The presence of vital pulp plays a pivotal role in root development. Therefore, when the pulp is reversibly inflamed it is crucial to maintain pulp vitality, especially in immature teeth (Deleimburg *et al.*, 2004). Caries and trauma are the main cause for pulp inflammation and necrosis. If these occur prior to root maturation, the root development would be halted, and can lead to an open apex (Harty, Parkins and Wengraf, 1970). Trauma to the anterior teeth is a relatively common occurrence during childhood especially in children of age 8-12 years, as this is a period of maximum physiologic growth and development and the children are actively involved in a lot of outdoor activities. Depending on the magnitude, trauma to maxillary anterior teeth may cause concussion, luxation, fracture, or avulsion of the teeth, in more severe cases, leads to necrosis of the pulp tissue (Moore, Howley and O'Connell, 2011). The maxillary anterior teeth tend to undergo many impact injuries because of its position in the jaw.

Pulp regeneration i.e. apexogenesis is not possible when long-term history of trauma causes external root resorption. In such cases, the treatment of choice is apexification or root-end closure. The endodontic treatment of immature permanent necrotic teeth is more difficult than conventional procedures because these teeth present widened root canals and open apices (Moore, Howley and O'Connell, 2011). The tooth roots may also suffer external infection-related (inflammatory) root resorption or alterations during treatment (Bakland and Andreasen, 2012). In such cases with infected pulps, it is necessary to use an intracanal dressing material to neutralize the bacteria and their products and to stimulate the apexification process by forming a mineralized apical barrier so that the subsequent condensation of gutta-percha can be properly achieved (Mohammadi and Dummer, 2011). Open apex complicates root canal treatment due to lack of apical stop, as there can be extravasation of irrigating solution and/or sealer into periradicular tissues, which can have a negative effect on the apical healing process. The main goal of treatment of teeth with pulpal necrosis is achieving an apical seal (Ghaziani, Aghasizadeh and Sheikh-Nezami, 2007) which is created by a barrier of hard tissue through a process known as apexification. (Floratos, Tsatsoulis and Kontakiotis, 2013).

Traditionally, apexification had been performed by using calcium hydroxide paste, due to its biological and healing performances (Yassen *et al.*, 2012). Calcium hydroxide has been successfully used for apical barrier formation in 74–100% of cases (Finucane and Kinirons, 1999). 86% of these treated teeth survived after a follow-up of 5 years. However, the use of calcium hydroxide paste use for apexification is not advocated these days as it involves a long treatment time and the prognosis is always uncertain (Hussainy *et al.*, 2018). The average length of time for apical barrier formation ranges from ~3 to 17 months, necessitating multiple visits for material replacement and delays the construction of the definitive restoration (Finucane and Kinirons, 1999). Exposure of the tissue to calcium hydroxide for long periods weakens the root structure, resulting in fractures, as well as induces periapical bone necrosis when there is overfilling of the material (Strom *et al.*, 2012). Another major disadvantage of apexification procedure using calcium hydroxide are the thin walls of the root which may fracture. Although the barrier is calcified it is actually porous and may contain a small amount of soft tissue (Manohar and Sharma, 2018; Teja and Ramesh, 2019).

As an alternative to traditional apexification using calcium hydroxide, a number of materials have been proposed in the literature (Ravinthar and Jayalakshmi, 2018; Teja, Ramesh and Priya, 2018). Among these materials, MTA is the most popular for open apex management (Janani, Palanivelu and Sandhya, 2020; Jose and Subbaiyan, 2020). MTA is composed of fine hydrophilic particles of tricalcium silicate, silicate oxide and tri calcium oxide. When mixed with sterile water it forms the colloidal gel and its setting time is about 3 to 4 hours in the presence of moisture (Nandakumar and Nasim, 2018). MTA has less leakage, better antibacterial properties, high marginal adaptation and short setting time of 4 hours, pH of 12.5 and is more biocompatible (Kubasad and Ghivari, 2011; Güneş and Aydinbelge, 2012).

With the use of mineral trioxide aggregate (MTA) in dentistry, it is possible to optimize the treatment time of open apex management by immediate placement of apical plug and the root canal filling (Ramamoorthi, Nivedhitha and Divyanand, 2015; R, Rajakeerthi and Ms, 2019). MTA has been shown to be a very effective root end filling material for sealing immature root canals with open apices that would otherwise impose technical challenges in obtaining adequate obturation (Kumar and Antony, 2018; Rajendran *et al.*, 2019). MTA has the ability to facilitate periradicular healing by inducing hard tissue formation (Camilleri and Pitt Ford, 2006). Mineral trioxide aggregate apart from being a root end filling material also has properties like regeneration of periradicular tissues like bone, cementum and periodontal ligament (Rajendran *et al.*, 2019; Siddique *et al.*, 2019). It has an excellent sealing ability, being a hydraulic cement material it sets even in the presence of moisture (Ramanathan and Solete, 2015; Noor, S Syed Shihaab and Pradeep, 2016).

Our department is passionate about research we have published numerous high quality articles in this domain over the past years (Abraham *et al.*, 2005; Devaki, Sathivel and BalajiRaghavendran, 2009; Neelakantan *et al.*, 2010, 2015; Arja *et al.*, 2013; Ramshankar *et al.*, 2014; Sumathi *et al.*, 2014; Surapaneni and Jainu, 2014; Surapaneni, Priya and Mallika, 2014; Ramamoorthi, Nivedhitha and Divyanand, 2015; Manivannan *et al.*, 2017; Ezhilarasan, 2018; Ezhilarasan, Sokal and Najimi, 2018; J *et al.*, 2018; Ravindiran and Praveenkumar, 2018; Malli Sureshbabu *et al.*, 2019; Mehta *et al.*, 2019; Krishnaswamy *et al.*, 2020; Samuel, Acharya and Rao, 2020; Sathish and Karthick, 2020)

This study was undertaken to analyse the number of open apex management done in maxillary incisors using MTA and its distribution and association with age and gender.

MATERIALS AND METHODS

Study Design and setting: In this cross sectional study, the data of 41 patients who underwent treatment for open apex management in Saveetha dental college were collected from dental records. Totally 47 maxillary incisors where open apex management was done using MTA were included in the study. At data extraction all information was anonymised and tabulated into a spreadsheet.

Ethical Approval: The study was commenced after approval from the institutional review board (Ethical approval number : SDC/SIHEC/2020/DIASDATA/0619-0320).

Selection of study population:

Inclusion criteria:

- Patients with necrotic pulp and open apex
- Patients without previous history for the treatment
- MTA used for apexification

Exclusion criteria:

- Patients who underwent incomplete treatment .
- Use of other materials for open apex management.

Sampling: Data collected from June 2019 to March 2020 comprised 41 patients who had undergone open apex management using MTA . The following data retrieved from the dental records: Patient's age, gender and tooth number .

Statistical analysis: The statistical analysis was done using SPSS software version 20.0 (SPSS Inc., Chicago, IL, USA). Descriptive statistics (percentage and mean) and Inferential statistics (Fisher's exact test) were done.

RESULTS AND DISCUSSION

Among the 47 teeth that underwent MTA apexification in our study, 82.98% belonged to males and 17.02% to females [Figure 1]. 51.06% of teeth belonged to patients <15 years of age and 48.94% belonged to those >15 years [Figure 2]. MTA apexification was done predominantly in males in both age groups. However, no significant association was found between gender and different age groups in patients that underwent MTA apexification (P value- 0.701 >0.05; Fisher's exact test).

Early necrosis of the pulp, incomplete root formation or external resorption of the root due to trauma results in blunted or shortened root with open apex (Pace, Giuliani and Pagavino, 2008; Araújo *et al.*, 2010). Apexification is the process of creating a hard tissue barrier at the root apex. Although calcium hydroxide was used most commonly for the process of apexification , the time duration for barrier formation was too long and the apical barrier formed was porous leading to reinfection (Ajwani and Saini, 2011). To overcome these disadvantages, MTA was introduced as a 'one visit Apexification material' (Stefopoulos, Tzanetakis and Kontakiotis, 2012).

MTA is one of the most effective materials for sealing the communication between endodontic and periodontal spaces (Torabinejad *et al.*, 1995; Al-Hezaimi *et al.*, 2005; Al-Kahtani *et al.*, 2005). MTA stimulates the production of interleukins and cytokines thereby promoting cementum like hard tissue formation, when in contact with the periradicular tissue. MTA plug in the apical portion of the root promotes apical repair and prevents root canal over-filling and increases the fracture resistance of immature teeth (Pace *et al.*, 2014).

Holland et al conducted a study and noticed biological closure of apical foramen and absence of inflammation in periapical tissue after the placement of MTA (Holland *et al.*, 2007) .Similarly , MTA has proven to be a successful material in acting as an apical barrier in many studies by eliminating the pre existing infection and promoting periodontal and bone healing.

(Güneş and Aydinbelge, 2012); (Raldi *et al.*, 2009); (Ajwani and Saini, 2011). In the study by Magro et al , treatment of open-apex incisors with placement of an apical matrix with lyophilized collagen sponge against which MTA apical plug can be condensed, had favorable prognosis . The radiographic assessment indicated local anatomical normality and total regression of the initial radiolucent lesion (Grazielle Magro *et al.*, 2017).

In a study done by Shrava et al , boys were found to have significantly higher numbers of tooth fractures than girls (Sharva *et al.*, 2017). Majority of previous studies also reported a higher frequency of dental trauma among boys than among girls, (Jesus *et al.*, 2010; Sulieman and Awooda, 2018; Carvalho and de Carvalho, 2019). This high incidence of trauma among boys could be a reason for a higher number of male patients seeking treatment for open apex as seen in our study [Figure 1]. A possible reason for male children being more prone to traumatic dental injuries than female children could be their participation and involvement in more aggressive sports and outdoor activities. The relatively low prevalence of trauma among females can also be explained by the fact that they are in general more mature in their behavior when compared with males, who tend to be more energetic and inclined toward exuberant outdoor activities.

The peak age to sustain traumatic dental injury was found to be in the age group of 13 to 14 years in a study conducted by Govindarajan et al (Govindarajan *et al.*, 2012). In our study, the number of teeth that underwent MTA apexification was slighter higher in the age group <15 years [Figure2]. No association was found between age and gender in teeth undergoing MTA apexification [Figure 3]. The effect of the patients' age on the outcome of apexification is controversial. It has been reported that apexification is indicated in young children (Lin, Chance and Skribner, 1986) and provides more favourable results, as shown in an animal study (Weinstein and Goldman, 1977). However, successful biologic apexification with Calcium hydroxide in adults has been reported in several case reports (Rotstein, Friedman and Katz, 1990; Gutmann and Fava, 1992; Calışkan and Türkün, 1997) . Clinical apexification treatment studies with MTA reported favourable results in immature teeth of different age groups ranging from 6 to 82 years, and composed of mostly children (Simon, Rilliard and Berdal, 2007; Holden *et al.*, 2008; Witherspoon *et al.*, 2008) .

The present study suggests that MTA apexification is done mostly in males and in the lesser age group suggesting people's willingness to address the problem immediately and preserve the natural tooth.

CONCLUSION

MTA apexification was mostly done in male population and in the age category less than 15 years. The placement of apical barrier using MTA is an alternative to conventional long-term calcium hydroxide which reduces the treatment time.

AUTHOR CONTRIBUTIONS

Priadarsini T, Dr Sowmya K were the main contributors for the concept, design, literature analysis, workshop discussions, drafting and revising the manuscript. Dr. Sowmya K and Dr. Dhanraj Ganapathy contributed to drafting and revising the manuscript. All authors gave final approval of the version to be published.

Conflicts of interest: There are no conflicts of interest.

REFERENCES

1. Abraham, S. *et al.* (2005) 'Evaluation of the inhibitory effect of triphala on PMN-type matrix metalloproteinase (MMP-9)', *Journal of periodontology*, 76(4), pp. 497–502.
2. Ajwani, P. and Saini, N. (2011) 'Non-surgical management of a mutilated maxillary central incisor with open apex and large periapical lesion', *Indian journal of dental research: official publication of Indian Society for Dental Research*, 22(3), pp. 475–477.
3. Al-Hezaimi, K. *et al.* (2005) 'Human saliva penetration of root canals obturated with two types of mineral trioxide aggregate cements', *Journal of endodontia*, 31(6), pp. 453–456.
4. Al-Kahtani, A. *et al.* (2005) 'In-vitro evaluation of microleakage of an orthograde apical plug of mineral trioxide aggregate in permanent teeth with simulated immature apices', *Journal of endodontia*, 31(2), pp. 117–119.
5. Araújo, R. A. *et al.* (2010) 'Single-session use of mineral trioxide aggregate as an apical barrier in a case of external root resorption', *Journal of oral science*, 52(2), pp. 325–328.
6. Arja, C. *et al.* (2013) 'Oxidative stress and antioxidant enzyme activity in South Indian male smokers with chronic obstructive pulmonary disease', *Respirology*, 18(7), pp. 1069–1075.
7. Bakland, L. K. and Andreasen, J. O. (2012) 'Will mineral trioxide aggregate replace calcium hydroxide in treating pulpal and periodontal healing complications subsequent to dental trauma? A review', *Dental traumatology: official publication of International Association for Dental Traumatology*, 28(1), pp. 25–32.
8. Calışkan, M. K. and Türkün, M. (1997) 'Periapical repair and apical closure of a pulpless tooth using calcium hydroxide', *Oral surgery, oral medicine, oral pathology, oral radiology, and endodontics*, 84(6), pp. 683–687.
9. Camilleri, J. and Pitt Ford, T. R. (2006) 'Mineral trioxide aggregate: a review of the constituents and biological properties of the material', *International endodontic journal*, 39(10), pp. 747–754.
10. Carvalho, R. T. da R. de and de Carvalho, R. T. da R. (2019) 'Prevalence and Factors Associated to Dental Erosion in 12-year-Old School Children from the City of Joaçaba (SC-Brazil)', *Journal of Dentistry and Oral Sciences*. doi: 10.37191/maps-ci-2582-3736-1(3)-018.
11. Deleimburg, M. *et al.* (2004) 'MTA Obturation of Pulpless Teeth with Open Apices: Bacterial Leakage as Detected by Polymerase Chain Reaction Assay', *Journal of Endodontics*, pp. 883–886. doi: 10.1097/01.don.0000128749.50151.24.
12. Devaki, T., Sathivel, A. and BalajiRaghavendran, H. R. (2009) 'Stabilization of mitochondrial and microsomal function by polysaccharide of *Ulva lactuca* on D-Galactosamine induced hepatitis in rats', *Chemico-biological interactions*, 177(2), pp. 83–88.
13. Ezhilarasan, D. (2018) 'Oxidative stress is bane in chronic liver diseases: Clinical and experimental perspective', *Arab journal of gastroenterology: the official publication of the Pan-Arab Association of Gastroenterology*, 19(2), pp. 56–64.
14. Ezhilarasan, D., Sokal, E. and Najimi, M. (2018) 'Hepatic fibrosis: It is time to go with hepatic stellate cell-specific therapeutic targets', *Hepatobiliary & pancreatic diseases international: HBPDI*, 17(3), pp. 192–197.
15. Finucane, D. and Kinirons, M. J. (1999) 'Non-vital immature permanent incisors: factors that may influence treatment outcome', *Dental traumatology: official publication of International Association for Dental Traumatology*, 15(6), pp. 273–277.
16. Floratos, S. G., Tsatsoulis, I. N. and Kontakiotis, E. G. (2013) 'Apical barrier formation after incomplete orthograde MTA apical plug placement in teeth with open apex--report of two cases', *Brazilian dental journal*, 24(2), pp. 163–166.
17. Ghaziani, P., Aghasizadeh, N. and Sheikh-Nezami, M. (2007) 'Endodontic treatment with MTA apical plugs: a case report', *Journal of oral science*, 49(4), pp. 325–329.
18. Govindarajan, M. *et al.* (2012) 'Prevalence of traumatic dental injuries to the anterior teeth among three to thirteen-year-old school children of Tamilnadu', *Contemporary clinical dentistry*, 3(2), pp. 164–167.
19. Grazielle Magro, M. *et al.* (2017) 'Endodontic Management of Open Apex Teeth Using Lyophilized Collagen Sponge and MTA Cement: Report of Two Cases', *Iranian endodontic journal*, 12(2), pp. 248–252.
20. Güneş, B. and Aydinbelge, H. A. (2012) 'Mineral trioxide aggregate apical plug method for the treatment of nonvital immature permanent maxillary incisors: Three case reports', *Journal of conservative dentistry: JCD*, 15(1), pp. 73–76.
21. Gutmann, J. L. and Fava, L. R. G. (1992) 'Periradicular healing and apical closure of a non-vital tooth in the presence of bacterial contamination', *International endodontic journal*, 25(6), pp. 307–311.

22. Harty, F. J., Parkins, B. J. and Wengraf, A. M. (1970) 'Success rate in root canal therapy. A retrospective study of conventional cases', *British dental journal*, 128(2), pp. 65–70.
23. Holden, D. T. *et al.* (2008) 'Clinical outcomes of artificial root-end barriers with mineral trioxide aggregate in teeth with immature apices', *Journal of endodontia*, 34(7), pp. 812–817.
24. Holland, R. *et al.* (2007) 'Influence of the type of vehicle and limit of obturation on apical and periapical tissue response in dogs' teeth after root canal filling with mineral trioxide aggregate', *Journal of endodontia*, 33(6), pp. 693–697.
25. Hussainy, S. N. *et al.* (2018) 'Clinical performance of resin-modified glass ionomer cement, flowable composite, and polyacid-modified resin composite in noncarious cervical lesions: One-year follow-up', *Journal of conservative dentistry: JCD*, 21(5), pp. 510–515.
26. Janani, K., Palanivelu, A. and Sandhya, R. (2020) 'Diagnostic accuracy of dental pulse oximeter with customized sensor holder, thermal test and electric pulp test for the evaluation of pulp vitality - An in vivo study', *Brazilian Dental Science*. doi: 10.14295/bds.2020.v23i1.1805.
27. Jesus, M. A. de *et al.* (2010) 'Epidemiologic survey of traumatic dental injuries in children seen at the Federal University of Rio de Janeiro, Brazil', *Brazilian oral research*, 24(1), pp. 89–94.
28. Jose, J. and Subbaiyan, H. (2020) 'Different Treatment Modalities followed by Dental Practitioners for Ellis Class 2 Fracture–A Questionnaire-based Survey', *The open dentistry journal*. Available at: <https://opendentistryjournal.com/VOLUME/14/PAGE/59/FULLTEXT/>.
29. J, P. C. *et al.* (2018) 'Prevalence and measurement of anterior loop of the mandibular canal using CBCT: A cross sectional study', *Clinical implant dentistry and related research*, 20(4), pp. 531–534.
30. Krishnaswamy, H. *et al.* (2020) 'Investigation of air conditioning temperature variation by modifying the structure of passenger car using computational fluid dynamics', *Thermal Science*, 24(1 Part B), pp. 495–498.
31. Kubasad, G. C. and Ghivari, S. B. (2011) 'Apexification with apical plug of MTA-report of cases', *Arch Oral Sci Res*, 1(2), pp. 104–107.
32. Kumar, D. and Antony, S. (2018) 'Calcified Canal and Negotiation-A Review', *Research Journal of Pharmacy and Technology*, 11(8), pp. 3727–3730.
33. Lin, L. M., Chance, K. and Skribner, J. (1986) 'Calcium hydroxide in endodontic therapy', *The Compendium of continuing education in dentistry*, 7(2), pp. 121, 126–30.
34. Malli Sureshababu, N. *et al.* (2019) 'Concentrated Growth Factors as an Ingenious Biomaterial in Regeneration of Bony Defects after Periapical Surgery: A Report of Two Cases', *Case reports in dentistry*, 2019, p. 7046203.
35. Manivannan, I. *et al.* (2017) 'Tribological and surface behavior of silicon carbide reinforced aluminum matrix nanocomposite', *Surfaces and Interfaces*, 8, pp. 127–136.
36. Manohar, M. P. and Sharma, S. (2018) 'A survey of the knowledge, attitude, and awareness about the principal choice of intracanal medicaments among the general dental practitioners and ...', *Indian journal of dental research: official publication of Indian Society for Dental Research*. Available at: <http://www.ijdr.in/article.asp?issn=0970-9290;year=2018;volume=29;issue=6;spage=716;epage=720;aulast=Manohar>.
37. Mehta, M. *et al.* (2019) 'Oligonucleotide therapy: An emerging focus area for drug delivery in chronic inflammatory respiratory diseases', *Chemico-biological interactions*, 308, pp. 206–215.
38. Mohammadi, Z. and Dummer, P. M. H. (2011) 'Properties and applications of calcium hydroxide in endodontics and dental traumatology', *International endodontic journal*, 44(8), pp. 697–730.
39. Moore, A., Howley, M. F. and O'Connell, A. C. (2011) 'Treatment of open apex teeth using two types of white mineral trioxide aggregate after initial dressing with calcium hydroxide in children', *Dental traumatology: official publication of International Association for Dental Traumatology*, 27(3), pp. 166–173.
40. Nandakumar, M. and Nasim, I. (2018) 'Comparative evaluation of grape seed and cranberry extracts in preventing enamel erosion: An optical emission spectrometric analysis', *Journal of conservative dentistry: JCD*, 21(5), pp. 516–520.
41. Neelakantan, P. *et al.* (2010) 'Root and Canal Morphology of Mandibular Second Molars in an Indian Population', *Journal of endodontics*, 36(8), pp. 1319–1322.
42. Neelakantan, P. *et al.* (2015) 'Photoactivation of curcumin and sodium hypochlorite to enhance antibiofilm efficacy in root canal dentin', *Photodiagnosis and photodynamic therapy*, 12(1), pp. 108–114.
43. Noor, S. S. S. E., S Syed Shihaab and Pradeep (2016) 'Chlorhexidine: Its properties and effects', *Research Journal of Pharmacy and Technology*, p. 1755. doi: 10.5958/0974-360x.2016.00353.x.
44. Pace, R. *et al.* (2014) 'Mineral trioxide aggregate as apical plug in teeth with necrotic pulp and immature apices: a 10-year case series', *Journal of endodontia*, 40(8), pp. 1250–1254.
45. Pace, R., Giuliani, V. and Pagavino, G. (2008) 'Mineral trioxide aggregate in the treatment of external invasive resorption: a case report', *International endodontic journal*, 41(3), pp. 258–266.
46. Rajendran, R. *et al.* (2019) 'Comparative Evaluation of Remineralizing Potential of a Paste Containing Bioactive Glass and a Topical Cream Containing Casein Phosphopeptide-Amorphous Calcium Phosphate:

- An in Vitro Study', *Pesquisa brasileira em odontopediatria e clinica integrada*, 19. Available at: http://www.scielo.br/scielo.php?pid=S1983-46322019000100364&script=sci_arttext.
47. Raldi, D. P. et al. (2009) 'Treatment options for teeth with open apices and apical periodontitis', *Journal*, 75(8), pp. 591–596.
 48. Ramamoorthi, S., Nivedhitha, M. S. and Divyanand, M. J. (2015) 'Comparative evaluation of postoperative pain after using endodontic needle and EndoActivator during root canal irrigation: A randomised controlled trial', *Australian endodontic journal: the journal of the Australian Society of Endodontology Inc*, 41(2), pp. 78–87.
 49. Ramanathan, S. and Solete, P. (2015) 'Cone-beam Computed Tomography Evaluation of Root Canal Preparation using Various Rotary Instruments: An in vitro Study', *The journal of contemporary dental practice*, 16(11), pp. 869–872.
 50. Ramshankar, V. et al. (2014) 'Risk stratification of early stage oral tongue cancers based on HPV status and p16 immunoeexpression', *Asian Pacific journal of cancer prevention: APJCP*, 15(19), pp. 8351–8359.
 51. Ravindiran, M. and Praveenkumar, C. (2018) 'Status review and the future prospects of CZTS based solar cell – A novel approach on the device structure and material modeling for CZTS based photovoltaic device', *Renewable and Sustainable Energy Reviews*, 94, pp. 317–329.
 52. Ravinthar, K. and Jayalakshmi (2018) 'Recent Advancements in Laminates and Veneers in Dentistry', *Research Journal of Pharmacy and Technology*, p. 785. doi: 10.5958/0974-360x.2018.00148.8.
 53. Rotstein, I., Friedman, S. and Katz, J. (1990) 'Apical closure of mature molar roots with the use of calcium hydroxide', *Oral surgery, oral medicine, and oral pathology*, 70(5), pp. 656–660.
 54. R, R., Rajakeerthi, R. and Ms, N. (2019) 'Natural Product as the Storage medium for an avulsed tooth – A Systematic Review', *Cumhuriyet Dental Journal*, pp. 249–256. doi: 10.7126/cumudj.525182.
 55. Samuel, S. R., Acharya, S. and Rao, J. C. (2020) 'School Interventions-based Prevention of Early-Childhood Caries among 3-5-year-old children from very low socioeconomic status: Two-year randomized trial', *Journal of public health dentistry*, 80(1), pp. 51–60.
 56. Sathish, T. and Karthick, S. (2020) 'Wear behaviour analysis on aluminium alloy 7050 with reinforced SiC through taguchi approach', *Journal of Materials Research and Technology*, 9(3), pp. 3481–3487.
 57. Sharva, V. et al. (2017) 'Traumatic dental injuries to the anterior teeth among 12-year and 15-year-old schoolchildren of urban and rural areas of Bhopal District, Central India: A prevalence study', *CHRISMED Journal of Health and Research*, 4(1), p. 38.
 58. Siddique, R. et al. (2019) 'Qualitative and quantitative analysis of precipitate formation following interaction of chlorhexidine with sodium hypochlorite, neem, and tulsi', *Journal of conservative dentistry: JCD*, 22(1), pp. 40–47.
 59. Simon, S., Rilliard, F. and Berdal, A. (2007) 'The use of mineral trioxide aggregate in one-visit apexification treatment: a prospective study', *International endodontic journal*. Available at: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-2591.2007.01214.x>.
 60. Stefopoulos, S., Tzanetakakis, G. N. and Kontakiotis, E. G. (2012) 'Non-surgical retreatment of a failed apicoectomy without retrofilling using white mineral trioxide aggregate as an apical barrier', *Brazilian dental journal*, 23(2), pp. 167–171.
 61. Strom, T. A. et al. (2012) 'Endodontic release system for apexification with calcium hydroxide microspheres', *Journal of dental research*, 91(11), pp. 1055–1059.
 62. Sulieman, A. G. and Awooda, E. M. (2018) 'Prevalence of Anterior Dental Trauma and Its Associated Factors among Preschool Children Aged 3–5 Years in Khartoum City, Sudan', *International journal of dentistry*, 2018. doi: 10.1155/2018/2135381.
 63. Sumathi, C. et al. (2014) 'Production of prodigiosin using tannery fleshing and evaluating its pharmacological effects', *TheScientificWorldJournal*, 2014, p. 290327.
 64. Surapaneni, K. M. and Jainu, M. (2014) 'Comparative effect of pioglitazone, quercetin and hydroxy citric acid on the status of lipid peroxidation and antioxidants in experimental non-alcoholic steatohepatitis', *Journal of physiology and pharmacology: an official journal of the Polish Physiological Society*, 65(1), pp. 67–74.
 65. Surapaneni, K. M., Priya, V. V. and Mallika, J. (2014) 'Pioglitazone, quercetin and hydroxy citric acid effect on cytochrome P450 2E1 (CYP2E1) enzyme levels in experimentally induced non alcoholic steatohepatitis (NASH)', *European review for medical and pharmacological sciences*, 18(18), pp. 2736–2741.
 66. Teja, K. V. and Ramesh, S. (2019) 'Shape optimal and clean more', *Saudi Endodontic Journal*. Available at: <http://www.saudiendodj.com/article.asp?issn=1658-5984;year=2019;volume=9;issue=3;spage=235;epage=236;aulast=Teja>.
 67. Teja, K. V., Ramesh, S. and Priya, V. (2018) 'Regulation of matrix metalloproteinase-3 gene expression in inflammation: A molecular study', *Journal of conservative dentistry: JCD*, 21(6), pp. 592–596.
 68. Torabinejad, M. et al. (1995) 'Physical and chemical properties of a new root-end filling material', *Journal of endodontia*, 21(7), pp. 349–353.

69. Weinstein, T. and Goldman, M. (1977) 'Apical hard-tissue deposition in adult teeth of monkeys with use of calcium hydroxide', *Oral surgery, oral medicine, and oral pathology*, 43(4), pp. 627–630.
70. Witherspoon, D. E. *et al.* (2008) 'Retrospective analysis of open apex teeth obturated with mineral trioxide aggregate', *Journal of endodontia*, 34(10), pp. 1171–1176.
71. Yassen, G. H. *et al.* (2012) 'The effect of frequency of calcium hydroxide dressing change and various pre- and inter-operative factors on the endodontic treatment of traumatized immature permanent incisors', *Dental Traumatology*, pp. 296–301. doi: 10.1111/j.1600-9657.2011.01089.x.

GRAPHS




Fig.1: Pie chart representing the distribution of teeth that underwent MTA apexification based on gender. 82.98% of the teeth that underwent MTA apexification belonged to males (Yellow) and 17.02% to females (Green).




Fig.2: Pie chart representing the distribution of teeth that underwent MTA apexification based on age group. 51.06% of the teeth that underwent MTA apexification belonged to the age group <15 years (Blue) and 48.04% to the group >15 years (Red).




Fig.3: Bar graph representing the association between age and gender in patients that underwent MTA apexification. X-axis represents the age groups and Y-axis represents the number of teeth involved with respect to gender . MTA apexification was done predominantly in males (Yellow) than females (Green) in both age groups. However, no significant association was found between gender and different age groups in patients that underwent MTA apexification (P value- 0.701 >0.05; Fisher’s exact test).