

---

## Incidence of Oral Mucosal Lesions in Patients Visiting A Private Dental Institution

---

FAHMIDA BINTI ABD RAHMAN<sup>1</sup>, VIVEK NARAYAN<sup>2\*</sup>, UMA MAHESHWARI<sup>3</sup>

<sup>1</sup>Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Science, Saveetha University, Chennai, India

<sup>2</sup>Senior Lecturer, Department of Oral Medicine and Radiology, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Science, Saveetha University, Chennai, India

<sup>3</sup>Department of Oral Medicine and Radiology, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Science, Saveetha University, Chennai 600077

\*Corresponding Author

Email ID: 151501005.sdc@saveetha.com<sup>1</sup>, viveknarayan@saveetha.com<sup>2</sup>, umamaheshwaritn@saveetha.com<sup>3</sup>

---

**Abstract:** Oral mucosal lesions act as a protective barrier against various harmful external agents. A wide variety of lesions and conditions either harmless or with serious complications may attack oral mucosa. Therefore, proper identification and treatment of these lesions play an important role in total oral health. It is a retrospective study. A total of 41439 patient records between January 2019 and December 2019 were evaluated in this study. Data such as age, gender, types of oral mucosal lesions were retrieved from the patient records of the private dental institution, Chennai. The data are then tabulated in Microsoft Excel and subsequently transferred to SPSS software for statistical analysis. Chi square test was done and the association between age and gender with types of oral mucosal lesions were found to be statistically significant. The incidence of oral mucosal lesions was 3.79 per 1000 per year. Overall, leukoplakia (27%) had the maximum predilection followed by smoker's palate (23%). Patients in the age group of 41 to 50 years were having the highest predilection for oral mucosal lesions (27.7%). Out of the 5 categories of oral mucosal lesions, potentially malignant disorders (60%) were highly prevalent in our study. This can be attributed to the increase in various adverse habits such as smoking, chewing tobacco, tobacco products and many more. More awareness about tobacco and its harmful effects must be promoted in order to prevent the rising numbers in potentially malignant disorders and oral cancer.

**Keywords:** Incidence; Oral mucosal lesions; Potentially malignant disorders; Smoking; Tobacco products, innovative

---

### INTRODUCTION

Oral health has an important role in the quality of life for an individual. Oral mucosa functions as an effective protective barrier and commonly attacked by pathologies that can be either harmful or harmless. Oral lesions may lead to impairment in speech and inability to eat (Dhanuthai *et al.*, 2016). Certain lesions may cause halitosis, dysesthesia or xerostomia. Factors such as deleterious habits, irregular or sharp teeth, ill-fitting prosthesis and poor oral hygiene can also contribute to the formation of oral mucosal lesions (Fleishman, Peles and Pisanti, 1985). Furthermore, other factors such as infection from bacteria, fungi, viruses, parasites, changes in immune system, systemic diseases, aging and neoplasia may also lead to oral mucosal lesions (Reichert, 2000). Epidemiologic studies impart information regarding prevalence, incidence and severity of oral disease in a particular population. It is essential to have the knowledge regarding distribution, etiology, risk factors and pathogenesis of oral mucosal lesions. This provides a timely primary prevention, early diagnosis and prompt treatment (Bhatnagar *et al.*, 2013). Several studies have documented the total range of possible lesions. The epidemiologic literature on oral mucosal diseases is still somewhat scanty even though the World Health Organization (WHO), "Guide to epidemiology and diagnosis of oral mucosal disease and conditions had been published in 1980. A study from Sankaranarayanan *et al.* (Sankaranarayanan *et al.*, 1997) showed that India has one of the highest rates of oral cancer varying from over 20 per 100,000 and less than 2 per 100,000 in the Middle East. Oral cancer accounts for almost 30% of all cancer in India. The diagnostic marker for oral cancer is by the expression of MMP-9 in the tissue (Venugopal and Uma Maheswari, 2016).

The prevalence of oral mucositis is usually gravitated to a type of cancer (Chaitanya *et al.*, 2017). Oral cancer can be caused by smokeless tobacco (Muthukrishnan and Warnakulasuriya, 2018). Oral hygiene is also important among the population to prevent any oral disease (Subashri and Uma Maheswari, 2016). Salivary markers are noninvasive diagnostic markers for the detection of malignant transformation of oral potentially

malignant disorders (OPMD) (Maheswari *et al.*, 2018). People who are exposed to an environmental background radiation such as ultraviolet rays may develop a PMD called actinic cheilosis (Muthukrishnan and Bijai Kumar, 2017). Malignancy is characterized by anaplasia, invasiveness, and metastasis (Misra *et al.*, 2015). Validated clinical competencies could provide a model for countries developing an advanced training curriculum for Oral Medicine (Steele *et al.*, 2015). Previously our team had conducted numerous studies which include radiographic study (Patil *et al.*, 2018), (Rohini and Jayanth Kumar, 2017), review studies (Chaitanya *et al.*, 2018), (Dharman and Muthukrishnan, 2016), (Subha and Arvind, 2019), clinical studies (Choudhury, 2015), (Muthukrishnan, Bijai Kumar and Ramalingam, 2016). Our department is passionate about research we have published numerous high quality articles in this domain over the past years ( (Kavitha *et al.*, 2014) , (Praveen *et al.*, 2001),(Devi and Gnanavel, 2014), (Putchala *et al.*, 2013), (Vijayakumar *et al.*, 2010), (Lekha *et al.*, 2014a, 2014b) (Danda, 2010) (Danda, 2010) (Parthasarathy *et al.*, 2016) (Gopalakannan, Senthilvelan and Ranganathan, 2012), (Rajendran *et al.*, 2019), (Govindaraju, Neelakantan and Gutmann, 2017), (P. Neelakantan *et al.*, 2015), (PradeepKumar *et al.*, 2016), (Sajan *et al.*, 2011), (Lekha *et al.*, 2014a), (Neelakantan, Grotra and Sharma, 2013), (Patil *et al.*, 2017), (Jeevanandan and Govindaraju, 2018), (Abdul Wahab *et al.*, 2017), (Eapen, Baig and Avinash, 2017), (Menon *et al.*, 2018), (Wahab *et al.*, 2018), (Vishnu Prasad *et al.*, 2018), (Uthrakumar *et al.*, 2010), (Ashok, Ajith and Sivanesan, 2017), (Prasanna Neelakantan *et al.*, 2015). Now we are focusing on retrospective studies, the aim of this study is to determine the incidence of oral mucosal lesions among patients attending a private dental institution in Tamil Nadu, Chennai.

## MATERIALS AND METHODS

This is a retrospective study. A total of 41439 case records of patients were evaluated between the period of January 2019 and December 2019. The study was conducted on patients between the ages less than 20 to 70 years and above. Case records which were complete and all types of oral mucosal lesions were included for the study. Incomplete case records were excluded. Convenient sampling method was used and photographic verification was done for cross verification of data. All the data was verified by a single trained examiner. The university set up of the study provided easy accessibility to data and provided a population with similar ethnicity. Ethical approval for this study was obtained from the institutional ethical committee (SDC/SIHEC/2020/DIASDATA/0619-0320). Data regarding age, gender, presence or absence of oral mucosal lesions were performed and the data was tabulated in Microsoft Excel. The data was then transferred to SPSS for statistical analysis. Independent variables included gender and age and dependent variables were the oral mucosal lesions. Chi square analysis was done to find out the association between variables.

## RESULTS AND DISCUSSION

In the present study, the total sample studied was 41439 case records of patients and among which males were 24051 and females were 17388. Among the 41439 patients 1647 of them had oral mucosal lesions. In the 1647 patients with oral mucosal lesions 1433 (3.46%) of them were males and 214 (0.52%) were females (figure 1). Chi square analysis was performed to find out the association between the gender of the patients and the oral mucosal lesions. The p value was found to be 0.00 which was less than 0.05 and this indicates the association between the gender and the presence of oral mucosal lesions is statistically significant (figure 1). This is similar to a study done by (Al-Maweri *et al.*, 2015) which reported that there was a significant difference ( $p < 0.05$ ) between males (80.3%) and females (69.6%). The incidence of the oral mucosal lesions in the present study was found to be 39.74 per 1000 per year or 3.97%. Males were more likely to have oral mucosal lesions compared to females. This finding is in accordance to studies from (Patil, Doni and Maheshwari, 2015), (Castellanos and Díaz-Guzmán, 2008), (Mehrotra *et al.*, 2010) and in contrast with (Al-Mobeeriek and AlDosari, 2009) which reported higher prevalence of oral mucosal lesions in females with a value of 57.7%, but study from (Corbet, Holmgren and Philipsen, 1994) reported that there was no difference in the prevalence between both gender. The higher prevalence in males in our current study could be due to higher numbers examined on males compared to females and due to higher prevalence risk habits among males than females.

The various oral mucosal lesions are categorised into 5 groups which are 1. Candida related oral lesions, 2. Potentially malignant disorders, 3. Reactive white lesions, 4. Pigmented lesions and 5. Ulcers and mucositis. Among these 5 groups, the most of the oral mucosal lesions were potentially malignant disorders (59.5%) which is followed by reactive white lesions (21.8%), ulcers and stomatitis (9.29%), candida related oral lesions (7.59%) and finally the least of the oral mucosal lesions were pigmented oral lesions (1.76%) (table 1). Among the potentially malignant disorders, leukoplakia had the highest predilection (27.81%) followed by smokers palate (22.59%), oral submucous fibrosis (OSMF) (6.86%), lichen planus (2.06%), erythroplakia (0.12%) and erythroleukoplakia (0.06%). This finding is similar to study from (Toum *et al.*, 2018) in which leukoplakia was the most common potentially malignant disorder (0.22%). In the category of reactive white lesions, tobacco pouch keratosis had the maximum predilection (14.94%) followed by frictional keratosis (5.22%) and chemical burn (1.64%). In ulcers and mucositis, traumatic ulcers had the highest presence (6.19%) followed by herpes labialis (2.98%) and mucositis (0.12%). Among the candida related lesions, denture stomatitis had the highest

occurrence (3.89%) followed by angular cheilitis (1.7%), oral thrush (1.46%) and erythematous candidiasis (0.55%). Finally the least among the oral mucosal lesions is the pigmented lesions and in that only smokers melanosis was present (1.76%) (table 1). Based on study from (Feng *et al.*, 2015) the highest prevalence for OMLs was fissured tongue (3.15%), followed by recurrent aphthous ulcer (1.48%), traumatic ulcer (1.13%) and angular cheilitis (0.86%). On comparing the results of the current study with the results of the study by Feng *et al.*, fissured tongue and recurrent aphthous ulcers were not seen in the present study and angular cheilitis was almost twice that of the value of Feng *et al.* and traumatic ulcer was almost 5 times that of the result of Feng *et al.* These variations can be due to the sample size of both the studies. The present study had a bigger sample size than that of Feng *et al.* and more number of oral mucosal lesions were observed in the present study.

In our study, certain types of oral mucosal lesions were more prevalent in a particular age group. Reactive white lesions were more prevalent in the age group of 20 years and less (0.61%). In all the other age groups potentially malignant disorders rank the highest (59.5%) among all the types of lesions. In contrast with findings from (Gambhir *et al.*, 2011) which reported that potentially malignant disorders like leukoplakia, lichen planus were the second most common findings in the study (22.2%). This variation in the ranking of the potentially malignant disorders in the study by Gambhir *et al.* and the current study could be because of the ethnicity and sample size. In the age groups 21 - 30 years, 31 - 40 years, 41 - 50 years and 51 - 60 years, the second highest was the reactive white lesions which were 6.74%, 6.5%, 5.34% and 1.88% respectively. Ulcers and mucositis were the second most common in the age groups 61 - 70 years and 70 years and greater and was found to be 1.03% and 0.12% respectively (figure 2). Chi square analysis was performed to find out the association between different age groups and different types of oral mucosal lesions. The p value was found to be 0.00 which was less than 0.05 and hence the association between age groups and the types of oral mucosal lesions are found to be statistically significant and oral mucosal lesions were more prevalent in the age group 41 - 50 years (figure 2). The finding is in contrast with a study from (Al-Maweri *et al.*, 2015) which revealed that the highest prevalence of oral mucosal lesions was in the age of 60-69 years old (80.8%). The increase in incidence of OMLs with increased age can be attributed to the fact that older people had long standing oral habits compared to younger populations and this variation with the current study could be due to ethnicity of the population of (Al-Maweri *et al.*, 2015).

Among the different genders, females had the highest predilection for candida related oral lesions (4.07%) followed by ulcers and mucositis (3.4%), potentially malignant disorders (3.28%) and reactive white lesions (2.25%). Males had the highest predilection for potentially malignant disorders (56.22%) followed by reactive white lesions (19.55%), ulcers and mucositis (5.95%), candida related oral lesions (3.52%) and the least occurrence was pigmented lesions (1.76%). Chi square analysis was done to find out the association between types of oral mucosal lesions and different genders. The p value was found to be 0.00 which is less than 0.05 and the association between types of oral mucosal lesions and gender was found to be statistically significant (figure 3). In the distribution of oral mucosal lesions based on gender, it can be observed that a study from (Gambhir *et al.*, 2011) reported carcinomas and potentially malignant disorders had the highest prevalence (80%) among males in comparison to all types of lesions. On comparing the results of their study and the present study, the present study did not have any malignancies but males had the predilection for potentially malignant disorders which was in accordance with the study done by (Gambhir *et al.*, 2011) and in their study females were more likely to be affected with soft tissue tumors and infections which was also partially in accordance with the present study where females were more affected for candida related oral lesions which is a fungal infection.

The present study had limitations such as the reliability of the findings and diagnosis present in the case records; this is because the findings are dependent on the skills of the oral diagnostician who made the initial clinical examination and diagnosis. This is again due to the retrospective type of the study design. All the lesions involved in the study are clinically diagnosed and no histopathological examination was performed. Future scope of the study can include a prospective and multicentric study design and patients habits and other etiological factors.

## CONCLUSION

Within the limit of our study, it can be concluded that male patients were commonly affected with oral mucosal lesions compared to female patients. The oral mucosal lesions were more prevalent in the 41 - 50 years age group and females had slightly higher occurrence of candida related oral lesions when compared to males. Among the various oral mucosal lesions, potentially malignant disorders were the highest in number.

## Author Contributions

Fahmida contributed towards data collection, data analysis, and manuscript preparation. Dr. Vivek Narayan contributed as a guide in performing this study.

## Conflict of Interest


There were no conflicts of interest declared by authors

## REFERENCES

1. Abdul Wahab, P. U. *et al.* (2017) 'Risk Factors for Post-operative Infection Following Single Piece Osteotomy', *Journal of maxillofacial and oral surgery*, 16(3), pp. 328–332.
2. Al-Maweri, S. *et al.* (2015) 'Oral mucosal lesions in elderly dental patients in Sana'a, Yemen', *Journal of International Society of Preventive and Community Dentistry*, p. 12. doi: 10.4103/2231-0762.156152.
3. Al-Mobeeriek, A. and AlDosari, A. M. (2009) 'Prevalence of oral lesions among Saudi dental patients', *Annals of Saudi Medicine*, pp. 365–368. doi: 10.4103/0256-4947.55166.
4. Ashok, B. S., Ajith, T. A. and Sivanesan, S. (2017) 'Hypoxia-inducible factors as neuroprotective agent in Alzheimer's disease', *Clinical and experimental pharmacology & physiology*, 44(3), pp. 327–334.
5. Bhatnagar, P. *et al.* (2013) 'Prevalence study of oral mucosal lesions, mucosal variants, and treatment required for patients reporting to a dental school in North India: In accordance with WHO guidelines', *Journal of family & community medicine*, 20(1), pp. 41–48.
6. Castellanos, J. L. and Díaz-Guzmán, L. (2008) 'Lesions of the oral mucosa: an epidemiological study of 23785 Mexican patients', *Oral surgery, oral medicine, oral pathology, oral radiology, and endodontics*, 105(1), pp. 79–85.
7. Chaitanya, N. *et al.* (2018) 'An insight and update on the analgesic properties of vitamin C', *Journal of Pharmacy And Bioallied Sciences*, p. 119. doi: 10.4103/jpbs.jpbs\_12\_18.
8. Chaitanya, N. C. *et al.* (2017) 'Role of Vitamin E and Vitamin A in Oral Mucositis Induced by Cancer Chemo/Radiotherapy- A Meta-analysis', *Journal of clinical and diagnostic research: JCDR*, 11(5), pp. ZE06–ZE09.
9. Choudhury, P. (2015) 'Vanishing Roots: First Case Report of Idiopathic Multiple Cervico–Apical External Root Resorption', *JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH*. doi: 10.7860/jcdr/2015/11698.5668.
10. Corbet, E. F., Holmgren, C. J. and Philipsen, H. P. (1994) 'Oral mucosal lesions in 65-74-year-old Hong Kong Chinese', *Community Dentistry and Oral Epidemiology*, pp. 392–395. doi: 10.1111/j.1600-0528.1994.tb01599.x.
11. Danda, A. K. (2010) 'Comparison of a single noncompression miniplate versus 2 noncompression miniplates in the treatment of mandibular angle fractures: a prospective, randomized clinical trial', *Journal of oral and maxillofacial surgery: official journal of the American Association of Oral and Maxillofacial Surgeons*, 68(7), pp. 1565–1567.
12. Devi, V. S. and Gnanavel, B. K. (2014) 'Properties of Concrete Manufactured Using Steel Slag', *Procedia Engineering*, 97, pp. 95–104.
13. Dhanuthai, K. *et al.* (2016) 'Geriatric oral lesions: A multicentric study', *Geriatrics & Gerontology International*, pp. 237–243. doi: 10.1111/ggi.12458.
14. Dharman, S. and Muthukrishnan, A. (2016) 'Oral mucous membrane pemphigoid - Two case reports with varied clinical presentation', *Journal of Indian Society of Periodontology*, 20(6), pp. 630–634.
15. Eapen, B. V., Baig, M. F. and Avinash, S. (2017) 'An Assessment of the Incidence of Prolonged Postoperative Bleeding After Dental Extraction Among Patients on Uninterrupted Low Dose Aspirin Therapy and to Evaluate the Need to Stop Such Medication Prior to Dental Extractions', *Journal of maxillofacial and oral surgery*, 16(1), pp. 48–52.
16. Feng, J. *et al.* (2015) 'Prevalence and distribution of oral mucosal lesions: a cross-sectional study in Shanghai, China', *Journal of Oral Pathology & Medicine*, pp. 490–494. doi: 10.1111/jop.12264.
17. Fleishman, R., Peles, D. B. and Pisanti, S. (1985) 'Oral mucosal lesions among elderly in Israel', *Journal of dental research*, 64(5), pp. 831–836.
18. Gambhir, R. S. *et al.* (2011) 'The prevalence of oral mucosal lesions in the patients visiting a dental school in Northern India in relation to sex, site and distribution: A retrospective study', *Journal of Clinical and Experimental Dentistry*, pp. e10–e17. doi: 10.4317/jced.3.e10.
19. Gopalakannan, S., Senthilvelan, T. and Ranganathan, S. (2012) 'Modeling and Optimization of EDM Process Parameters on Machining of Al 7075-B4C MMC Using RSM', *Procedia Engineering*, 38, pp. 685–690.
20. Govindaraju, L., Neelakantan, P. and Gutmann, J. L. (2017) 'Effect of root canal irrigating solutions on the compressive strength of tricalcium silicate cements', *Clinical oral investigations*, 21(2), pp. 567–571.
21. Jeevanandan, G. and Govindaraju, L. (2018) 'Clinical comparison of Kedo-S paediatric rotary files vs manual instrumentation for root canal preparation in primary molars: a double blinded randomised clinical trial', *European Archives of Paediatric Dentistry*, pp. 273–278. doi: 10.1007/s40368-018-0356-6.
22. Kavitha, M. *et al.* (2014) 'Solution combustion synthesis and characterization of strontium substituted hydroxyapatite nanocrystals', *Powder Technology*, 253, pp. 129–137.
23. Lekha, L. *et al.* (2014a) 'Schiff base complexes of rare earth metal ions: Synthesis, characterization and

- catalytic activity for the oxidation of aniline and substituted anilines', *Journal of organometallic chemistry*, 753, pp. 72–80.
24. Lekha, L. *et al.* (2014b) 'Synthesis, spectroscopic characterization and antibacterial studies of lanthanide(III) Schiff base complexes containing N, O donor atoms', *Journal of Molecular Structure*, pp. 307–313. doi: 10.1016/j.molstruc.2013.10.014.
  25. Maheswari, T. N. U. *et al.* (2018) 'Salivary micro RNA as a potential biomarker in oral potentially malignant disorders: A systematic review', *Tzu Chi Medical Journal*, p. 55. doi: 10.4103/tcmj.tcmj\_114\_17.
  26. Mehrotra, R. *et al.* (2010) 'Prevalence of oral soft tissue lesions in Vidisha', *BMC Research Notes*, p. 23. doi: 10.1186/1756-0500-3-23.
  27. Menon, S. *et al.* (2018) 'Selenium nanoparticles: A potent chemotherapeutic agent and an elucidation of its mechanism', *Colloids and surfaces. B, Biointerfaces*, 170, pp. 280–292.
  28. Misra, S. R. *et al.* (2015) 'Metastatic hepatocellular carcinoma in the maxilla and mandible, an extremely rare presentation', *Contemporary clinical dentistry*, 6(Suppl 1), pp. S117–21.
  29. Muthukrishnan, A. and Bijai Kumar, L. (2017) 'Actinic cheilosis: early intervention prevents malignant transformation', *BMJ case reports*, 2017. doi: 10.1136/bcr-2016-218654.
  30. Muthukrishnan, A., Bijai Kumar, L. and Ramalingam, G. (2016) 'Medication-related osteonecrosis of the jaw: a dentist's nightmare', *BMJ case reports*, 2016. doi: 10.1136/bcr-2016-214626.
  31. Muthukrishnan, A. and Warnakulasuriya, S. (2018) 'Oral health consequences of smokeless tobacco use', *The Indian journal of medical research*, 148(1), pp. 35–40.
  32. Neelakantan, P. *et al.* (2015) 'Antibiofilm activity of three irrigation protocols activated by ultrasonic, diode laser or Er:YAG laser in vitro', *International endodontic journal*, 48(6), pp. 602–610.
  33. Neelakantan, P. *et al.* (2015) 'Influence of Irrigation Sequence on the Adhesion of Root Canal Sealers to Dentin: A Fourier Transform Infrared Spectroscopy and Push-out Bond Strength Analysis', *Journal of endodontia*, 41(7), pp. 1108–1111.
  34. Neelakantan, P., Grotra, D. and Sharma, S. (2013) 'Retreatability of 2 mineral trioxide aggregate-based root canal sealers: a cone-beam computed tomography analysis', *Journal of endodontia*, 39(7), pp. 893–896.
  35. Parthasarathy, M. *et al.* (2016) 'Effect of hydrogen on ethanol-biodiesel blend on performance and emission characteristics of a direct injection diesel engine', *Ecotoxicology and environmental safety*, 134(Pt 2), pp. 433–439.
  36. Patil, S. B. *et al.* (2017) 'Comparison of Extended Nasolabial Flap Versus Buccal Fat Pad Graft in the Surgical Management of Oral Submucous Fibrosis: A Prospective Pilot Study', *Journal of maxillofacial and oral surgery*, 16(3), pp. 312–321.
  37. Patil, S., Doni, B. and Maheshwari, S. (2015) 'Prevalence and Distribution of Oral Mucosal Lesions in a Geriatric Indian Population', *Canadian Geriatrics Journal*. doi: 10.5770/cgj.18.123.
  38. Patil, S. R. *et al.* (2018) 'Three-Rooted Mandibular First Molars in a Saudi Arabian Population: A CBCT Study', *Pesquisa Brasileira em Odontopediatria e Clínica Integrada*, p. e4133. doi: 10.4034/pboci.2018.181.87.
  39. PradeepKumar, A. R. *et al.* (2016) 'Diagnosis of Vertical Root Fractures in Restored Endodontically Treated Teeth: A Time-dependent Retrospective Cohort Study', *Journal of endodontia*, 42(8), pp. 1175–1180.
  40. Praveen, K. *et al.* (2001) 'Hypotensive anaesthesia and blood loss in orthognathic surgery: a clinical study', *The British journal of oral & maxillofacial surgery*, 39(2), pp. 138–140.
  41. Putchala, M. C. *et al.* (2013) 'Ascorbic acid and its pro-oxidant activity as a therapy for tumours of oral cavity – A systematic review', *Archives of Oral Biology*, pp. 563–574. doi: 10.1016/j.archoralbio.2013.01.016.
  42. Rajendran, R. *et al.* (2019) 'Comparative Evaluation of Remineralizing Potential of a Paste Containing Bioactive Glass and a Topical Cream Containing Casein Phosphopeptide-Amorphous Calcium Phosphate: An in Vitro Study', *Pesquisa Brasileira em Odontopediatria e Clínica Integrada*, pp. 1–10. doi: 10.4034/pboci.2019.191.61.
  43. Reichart, P. A. (2000) 'Oral mucosal lesions in a representative cross-sectional study of aging Germans', *Community Dentistry and Oral Epidemiology*, pp. 390–398. doi: 10.1034/j.1600-0528.2000.028005390.x.
  44. Rohini, S. and Jayanth Kumar, V. (2017) 'Incidence of dental caries and pericoronitis associated with impacted mandibular third molar-A radiographic study', *Research Journal of Pharmacy and Technology*, p. 1081. doi: 10.5958/0974-360x.2017.00196.2.
  45. Sajan, D. *et al.* (2011) 'Molecular structure and vibrational spectra of 2,6-bis(benzylidene)cyclohexanone: a density functional theoretical study', *Spectrochimica acta. Part A, Molecular and biomolecular spectroscopy*, 78(1), pp. 113–121.
  46. Sankaranarayanan, R. *et al.* (1997) 'Chemoprevention of oral leukoplakia with vitamin A and beta carotene: an assessment', *Oral Oncology*, pp. 231–236. doi: 10.1016/s0964-1955(97)00010-9.

47. Steele, J. C. *et al.* (2015) ‘World Workshop on Oral Medicine VI: an international validation study of clinical competencies for advanced training in oral medicine’, *Oral surgery, oral medicine, oral pathology and oral radiology*, 120(2), pp. 143–51.e7.
48. Subashri, A. and Uma Maheshwari, T. N. (2016) ‘Knowledge and attitude of oral hygiene practice among dental students’, *Research Journal of Pharmacy and Technology*, p. 1840. doi: 10.5958/0974-360x.2016.00375.9.
49. Subha, M. and Arvind, M. (2019) ‘Role of Magnetic Resonance Imaging in Evaluation of Trigeminal Neuralgia with its Anatomical Correlation’, *Biomedical and Pharmacology Journal*, pp. 289–296. doi: 10.13005/bpj/1640.
50. Toum, S. E. *et al.* (2018) ‘Prevalence and Distribution of Oral Mucosal Lesions by Sex and Age Categories: A Retrospective Study of Patients Attending Lebanese School of Dentistry’, *International Journal of Dentistry*, pp. 1–6. doi: 10.1155/2018/4030134.
51. Uthrakumar, R. *et al.* (2010) ‘Bulk crystal growth and characterization of non-linear optical bithiourea zinc chloride single crystal by unidirectional growth method’, *Current applied physics: the official journal of the Korean Physical Society*, 10(2), pp. 548–552.
52. Venugopal, A. and Uma Maheshwari, T. N. (2016) ‘Expression of matrix metalloproteinase-9 in oral potentially malignant disorders: A systematic review’, *Journal of oral and maxillofacial pathology: JOMFP*, 20(3), pp. 474–479.
53. Vijayakumar, G. N. S. *et al.* (2010) ‘Synthesis of electrospun ZnO/CuO nanocomposite fibers and their dielectric and non-linear optic studies’, *Journal of alloys and compounds*, 507(1), pp. 225–229.
54. Vishnu Prasad, S. *et al.* (2018) ‘Report on oral health status and treatment needs of 5-15 years old children with sensory deficits in Chennai, India’, *Special care in dentistry: official publication of the American Association of Hospital Dentists, the Academy of Dentistry for the Handicapped, and the American Society for Geriatric Dentistry*, 38(1), pp. 58–59.
55. Wahab, P. U. A. *et al.* (2018) ‘Scalpel Versus Diathermy in Wound Healing After Mucosal Incisions: A Split-Mouth Study’, *Journal of oral and maxillofacial surgery: official journal of the American Association of Oral and Maxillofacial Surgeons*, 76(6), pp. 1160–1164.




**Fig.1:** represents the association of gender of the patients with presence (black) and absence (orange) of oral mucosal lesions and the X axis represents gender of the patients and Y axis represents the number of patients. Chi square test was done and the association was found to be statistically significant as the p value= 0.00 (<0.05). Male patients were commonly affected with oral mucosal lesions compared to female patients.


**Table 1:** depicts the different categories of oral mucosal lesions and the total number of the types of oral mucosal lesions in each category. The table reveals that potentially malignant disorders (59.5%) are the highest followed by reactive white lesions (21.8%), ulcers and mucositis (9.29%), candida related oral lesions (7.59%) and pigmented oral lesions (1.76%).

Category	Oral lesions	Number, (%)	Total, (%)
----------	--------------	-------------	------------

Candida related oral lesions	Angular cheilitis	28 (1.70%)	125(7.59%)
	Oral thrush	24 (1.46%)	
	Denture stomatitis	64 (3.89%)	
	Erythematous candidiasis	9 (0.55%)	
Potentially malignant disorders	Leukoplakia	458 (27.81%)	980(59.50%)
	Lichen planus	34 (2.06%)	
	OSMF	113 (6.86%)	
	Smokers palate	372 (22.59%)	
	Erythroleukoplakia	1 (0.06%)	
	Erythroplakia	2 (0.12%)	
Reactive white lesions	Chemical burn	27 (1.64%)	359(21.80%)
	Frictional keratosis	86 (5.22%)	
	Tobacco pouch keratosis	246 (14.94%)	
Pigmented lesions	Smokers melanosis	29 (1.76%)	29(1.76%)
Ulcers and mucositis	Traumatic ulcer	102 (6.19%)	153(9.29%)
	Mucositis	2 (0.12%)	
	Herpes labialis	49 (2.98%)	
Total		1647(100%)	



**Fig.2:** represents association between age groups and types of oral mucosal lesions like candida related oral lesions (blue), potentially malignant disorders (green), reactive white lesions (light green), pigmented lesions (violet) and ulcer and mucositis (yellow). The X axis represents the age and Y axis represents the number of patients having oral mucosal lesions. Chi square test was done and the association was found to be statistically significant with p value =0.00 (<0.05). The oral mucosal lesions were more prevalent in the 41 - 50 years age group.



**Fig.3:** represents association between gender and types of oral mucosal lesions and the X axis represents gender of the patients and Y axis represents the number of patients having oral mucosal lesions. Chi square test was done and the association was found to be statistically significant with p value =0.00 (<0.05). Males had the highest occurrence of potentially malignant lesions, reactive white lesions, pigmented lesions and ulcers and mucositis when compared to females. Females had slightly higher occurrence of candida related oral lesions when compared to males.